

Simultaneous two-dimensional best Diophantine approximations in the Euclidean norm

Evgeny V. Ermakov

1. Introduction

This paper is devoted to the exponents of growth of denominators of best simultaneous Diophantine approximations. Consider $\mathbb{R}^n$ with a norm $\|\cdot\|$. For any vector $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n \setminus \mathbb{Q}^n$ and any $q \in \mathbb{Z}$ define the following value:

$$\delta_q = \min_{\mathbf{p}=(p_1, \dots, p_n) \in \mathbb{Z}^n} \|q \cdot \alpha - \mathbf{p}\|.$$

Let $\mathbf{p}(q) \in \mathbb{Z}^n$ be the vector, where the minimum is attained; let $\mathbf{r}(q) = q \cdot \alpha - \mathbf{p}(q)$, so $\delta_q = \|\mathbf{r}(q)\|$. Given a norm $\|\cdot\|$ in $\mathbb{R}^n$ and a vector $\alpha \in \mathbb{R}^n \setminus \mathbb{Q}^n$ we can define the sequence of best approximations (with respect to this norm) as a sequence $(q_k)_{k=1}^\infty$, such that $q_1 = 1$ and $\forall q < q_k \quad \delta_q > \delta_{q_k}$. Now we can define following values:

$$g(\alpha, \|\cdot\|) = \liminf_{k \rightarrow \infty} (q_k)^{\frac{1}{k}},$$

$$G(n, \|\cdot\|) = \inf_{\alpha \in \mathbb{R}^n \setminus \mathbb{Q}^n} g(\alpha, \|\cdot\|).$$

J.Lagarias [1] has proved the following statement:

Theorem 1. *For any norm $\|\cdot\|$ on $\mathbb{R}^n$ and a vector α , that has at least one irrational coordinate, the inequality $q_{k+2^{n+1}} \geq 2q_{k+1} + q_k$ holds for all $k \geq 1$. So $G(n, \|\cdot\|) \geq \theta$, where θ is the maximal positive root of $\theta^{2^{n+1}} = 2\theta + 1$.*

In this paper we consider $\mathbb{R}^2$ with the Euclidian norm. From Theorem 1 it follows that for the Euclidian norm in $\mathbb{R}^2$, and any vector α one has $q_{k+8} \geq q_{k+1} + q_k$.

There is another well known statement that holds for any norm. Given a norm $\|\cdot\|$ in $\mathbb{R}^n$ consider the contact number $K(n, \|\cdot\|)$. This number is defined as the maximal number of unit balls with respect to the norm $\|\cdot\|$ without interior common points that can touch another unit ball.

Theorem 2. *For any norm $\|\cdot\|$ on $\mathbb{R}^n$ with the contact number $K = K(n, \|\cdot\|)$ and a vector α , that has at least one irrational coordinate, we have the inequality $q_{k+K} \geq q_{k+1} + q_k$, and so $G(\|\cdot\|) \geq \theta$, where θ is maximum positive root of $\theta^K = \theta + 1$.*

For the Euclidian norm in $\mathbb{R}^2$ we have $K = 6$. So Theorem 2 gives the inequality

$$q_{k+6} \geq q_{k+1} + q_k. \tag{1}$$

It follows that $G(2, \|\cdot\|_e) \geq \theta$, where θ is maximum positive root of $\theta^6 = \theta + 1$ and $\|\cdot\|_e$ is the Euclidian norm.

Theorem 2 is a well known result, one can find a proof of it in M.Romanov paper [3]. M.Romanov [3] proved a stronger result that the inequality

$$q_{k+4} \geq q_{k+1} + q_k. \quad (2)$$

is valid for any $k \geq 1$. From inequality (2) it follows that $G(2, \|\cdot\|_e) \geq \theta_0$ where θ_0 is a positive root of $\theta_0^6 = \theta_0 + 1$, $\theta_0 = 1.220744$. The main result of the present paper is an improvement of Romanov's result.

Theorem 3. *For the Euclidian norm in $\mathbb{R}^2$ and any vector α , that has at least one irrational coordinate one has $G(2, \|\cdot\|_e) \geq 1.228043$.*

The proof of Theorem 3 is based on following geometric statement that together with the inequality (2) and some numerical calculations gives the lower bound.

Theorem 4. *Suppose that $\alpha \in \mathbb{R}^2$ has at least one irrational coordinate. Let $q_k \dots q_{k+4}$ be consecutive denominators from the sequence of best approximations in Euclidian norm for vector α . Then for every $k \geq 1$ at least one of two following inequalities are valid:*

$$q_{k+3} + q_{k+2} \geq 2q_{k+1} + q_k \quad (3)$$

$$q_{k+4} \geq q_{k+2} + q_k \quad (4)$$

Moreover, among any two successive values of k for at least one value the inequality (3) holds.

A.Brentjes [2] gave the following example. Let η be the maximal root of the equation $\eta^3 = \eta + 1$, $\eta = 1.3248\dots$. Then for $\alpha = (\alpha_1, \alpha_2) = (\eta, \eta^2)$ one has $g(\alpha, \|\cdot\|_e) = \eta$. J.Lagarias [1] made a conjecture, that $G(2, (\|\cdot\|_e)) = \eta$.

In Sections 2,3 below we give a complete proof of Theorem 4. In Section 4 we deduce Theorem 3 from Theorem 4 and Romanov's theorem. There we describe all necessary computer calculations.

2. Geometric lemmas

Lemma 1. *Consider a convex hexagon $A_1A_2A_3A_4A_5A_6$. Suppose that its opposite sides are equal and parallel. Suppose that O is an interior point of the hexagon. Let all the distances $|A_1O|, |A_2O|, |A_3O|, |A_5O|$ are different. Then there exists $i \in \{1, 2, 3, 5\}$ such that*

$$|A_iO| > \min_{j=1,2,3} |A_jA_{j+1}|.$$

Proof. Let $a = \min(|A_1A_2|, |A_2A_3|)$. Without loss of generality suppose that $a = |A_1A_2|$.

Consider circles ω_1 and ω_2 with radiuses a and centers in A_1 and A_3 correspondingly. Let κ_1 and κ_2 be closed disks bounded by ω_1 and ω_2 . Define $\Omega = \kappa_1 \cap \kappa_2$. (See fig.1.)

Suppose that the conclusion of Lemma 1 is not true, that is there exists an interior point O of hexagon $A_1A_2A_3A_4A_5A_6$ such that $|A_1O|, |A_2O|, |A_3O|, |A_5O|$ are different and

$$|A_iO| \leq \min_{j=1,2,3} |A_jA_{j+1}|, \quad i = 1, 2, 3, 5.$$

self intersections.

- Lemma 2.** 1. The tetragon $R_1R_2R_3R_4$ is convex, point O lies inside it.
 2. All of its sides and diagonals are not less than the longest remainder vector $|r(q_k)|$.
 3. Angles between vectors $\overrightarrow{OR_i}$ and $\overrightarrow{OR_j}$ ($i \neq j$) are greater than $\frac{\pi}{3}$.

Proof. Suppose, that $|R_iR_j| < |r(q_k)|$ for any $i \neq j$. Let R_i, R_j are the endpoints of vectors $\mathbf{r}(q_s)$ and $\mathbf{r}(q_l)$ correspondingly. Then $|\mathbf{r}(q_s - q_l)| < |\mathbf{r}(q_k)|$. From $q_{k+3} < q_{k+1} + q_k$ it follows that $0 < |q_s - q_l| < q_{k+1}$. Last inequalities contradict to the fact that q_k and q_{k+1} are denominators of consecutive best approximations. The second statement of Lemma 2 is proved.

In any triangle $OR_iR_j, i \neq j$ the side R_iR_j is the greatest one. Lengths of $\overrightarrow{\mathbf{r}(q_k)}$ decrease strictly, so those triangles can not have three equal sides and angles between vectors $\overrightarrow{OR_i}$ are greater then $\frac{\pi}{3}$. Other angles in these triangles are less or equal to $\frac{\pi}{3}$. We see that $R_1R_2R_3R_4$ is convex, and the point O lies inside it. Lemma 2 is proved.

3. Proof of Theorem 4

We need two more lemmas.

- Lemma 3.** If tetragon $R_1R_2R_3R_4$ is not a parallelogram, then the inequality (3) holds.

Proof. If $R_1R_2R_3R_4$ has no parallel sides, then we can make a convex hexagon by building parallelograms on two pairs of its sides. (See fig.2.) Without loss of generality we may suppose that the hexagon vertex R_4 lies between the vertices X_1 and X_2 . So we have constructed the hexagon $R_1R_2R_3X_2R_4X_1$.




fig.2

Consider the segment R_3X_1 (it is equal and parallel to segment R_1X_2). Put $x = |R_3X_1|$. By the construction the length of the remainder vector for the denominator $q = |q_1 + q_3 - q_2 - q_4|$ is not greater than x .

As the sequence (q_k) increases strictly, we have three possible values of q . So we should consider three cases.

Case 1. $q = |q_{k+3} + q_k - q_{k+2} - q_{k+1}|$. Here $0 < q < q_k$, and the length of the remainder vector for q is not less than $|\mathbf{r}(q_{k-1})|$. So $x \geq |\mathbf{r}(q_k)|$.

Case 2. $q = q_{k+3} + q_{k+1} - q_{k+2} - q_k$. Here $0 < q < q_{k+1}$. The length of the remainder vector for q is not less than $|\mathbf{r}(q_k)|$ (q is the denominator of the next best approximation). So $x \geq |\mathbf{r}(q_k)|$.

Case 3. $q = q_{k+3} + q_{k+2} - q_{k+1} - q_k$. Then $q > 0$ and we have 2 subcases:

3a $q = q_{k+3} + q_{k+2} - q_{k+1} - q_k < q_{k+1}$. Here as in cases 1 and 2 we have $x \geq |\mathbf{r}(q_k)|$.

3b $q = q_{k+3} + q_{k+2} - q_{k+1} - q_k \geq q_{k+1}$ Here we get the inequality (3).

In cases **1**, **2**, **3a** we have the following situation. As $|\mathbf{r}(q_k)| > |\mathbf{r}(q_{k+1})| > |\mathbf{r}(q_{k+2})| > |\mathbf{r}(q_{k+3})|$ we see that the hexagon $R_1R_2R_3X_2R_4X_1$ and the zero point O satisfy the conditions of Lemma 1. By Lemma 1 we see that

$$\max_{i=1,2,3,4} |R_iO| > \min\{|X_1R_1|, |R_1R_2|, |R_2R_3|\}.$$

As in our cases $x \geq |\mathbf{r}(q_k)|$ we see that

$$\max_{i=1,2,3,4} |R_iO| > \min\{|R_4R_1|, |R_1R_2|, |R_2R_3|, |R_3R_4|\}.$$

This contradicts to Lemma 2. So the cases are **1**, **2**, **3a** are not possible.

But in the remaining case **3b** we have the inequality (3).

To finish the proof of Lemma 3 we must consider the case when $R_1R_2R_3R_4$ has a pair of parallel sides. Then the hexagon $R_1R_2R_3X_2R_4X_1$ is a degenerate one (two its angles are equal to π). Now the proof follows the steps of the proof in non-degenerate case. The only difference is that we apply Lemma 1 for the degenerate hexagon. Lemma 3 is proved.

Lemma 4. *If $R_1R_2R_3R_4$ is a parallelogram and $q_{k+3} < q_{k+1} + q_k$, then endpoints of the next four remainder vectors (for $k+1, k+2, k+3, k+4$) do not form a parallelogram.*

Proof. Suppose they do. Let $\mathbf{r}(q_{k+4}) = \overline{OR_5}$, $\mathbf{r}(q_k) = \overline{OR}$. This parallelogram has three common vertices with $R_1R_2R_3R_4$. So one of the vertices of the hexagon $R_1R_2R_3R_4$ is the center of the segment RR_5 . This vertex we denote by R_6 .

As $\|OR\| = |\mathbf{r}(q_k)| > |\mathbf{r}(q_{k+4})| = \|OR_5\|$, we see that the zero point O lies closer to R_5 than to R . So in the triangle ORR_6 the angle in the vertex R_6 is greater than $\frac{\pi}{3}$ and the length of the remainder vector $\mathbf{r}(q_k) = \overline{OR}$ is greater than the length of the parallelogram's side RR_2 . We get the contradiction to Lemma 2. Lemma 4 is proved.

Proof of Theorem 4.

1. If points R_1, R_2, R_3, R_4 do not form a parallelogram, then using Lemma 3 we get inequality (3).

2. If the inequality $q_{k+3} < q_{k+1} + q_k$ do not holds, we again get inequality (3).

3. We may suppose that R_1, R_2, R_3, R_4 do form a parallelogram and $q_{k+3} < q_{k+1} + q_k$. Then by Lemma 4 the endpoints of the next four remainder vectors (for $k+1, k+2, k+3, k+4$) do not form a parallelogram. So for the approximations $k+1, k+2, k+3, k+4$ the inequality (3) is valid. We see that

$$q_{k+4} + q_{k+3} \geq 2q_{k+2} + q_{k+1}. \quad (5)$$

Let the endpoints of vectors

$$\mathbf{r}(q_k), \mathbf{r}(q_{k+1}), \mathbf{r}(q_{k+2}), \mathbf{r}(q_{k+3})$$

form a parallelogram in the order

$$\mathbf{r}(\hat{q}_1), \mathbf{r}(\hat{q}_2), \mathbf{r}(\hat{q}_3), \mathbf{r}(\hat{q}_4).$$

Then the remainder vector for the denominator $p = |\hat{q}_1 + \hat{q}_3 - \hat{q}_2 - \hat{q}_4|$ is equal to zero. As α is not a rational vector we see that $p = 0$. As the sequence of denominators of best approximations increases strictly we get $0 = p = q_k + q_{k+3} - q_{k+2} - q_{k+1}$.

The last equality together with (5) implies (8). Theorem 4 is proved.

4. Proof of Theorem 3

From Theorem 4 we immediately obtain

Proposition 1. Let $l \in \mathbb{R}$. Let $\alpha \in \mathbb{R}^2 \setminus \mathbb{Q}^2$. Let $\epsilon \in \mathbb{R}$. Then for every $k \geq 1$ for five consecutive denominators $q_k, \dots, q_{k+4}$ we have at least one of three following inequalities

$$q_{k+2} \geq lq_{k+1} \quad (6)$$

$$q_{k+3} \geq (2-l)q_{k+1} + q_k \quad (7)$$

$$q_{k+4} \geq q_{k+2} + q_k \quad (8)$$

Moreover, for any two successive values of k for at least one value the inequality (6) or the inequality (7) holds.

For further proof we need to use some computer calculations.

Let $0 < l < 2$. Let $m = 1, \dots, 7$.

Put $r_0 = r_1 = 31$, $r_2 = r_3 = r_4 = 33$, $r_5 = 34$, $r_6 = 35$,

$l_0 = \dots = l_3 = 1.298$, $l_4 = l_5 = l_6 = 1.293$,

$\theta_0 = 1.2207$, $\theta_1 = 1.2272$, $\theta_2 = 1.2275$, $\theta_3 = 1.22779$,

$\theta_4 = 1.2278$, $\theta_5 = 1.22785$, $\theta_6 = 1.22791$, $\theta_7 = 1.228043$.

Consider a sequence $I = (i_0, \dots, i_{r-1})$, $i_\nu \in \{1, 2, 3\}$ such that in any couple $i_\nu, i_{\nu+1}$ at least one element is not equal to 3. For such I we construct a sequence $\{Q_k(I, m)\}$, $0 \leq k \leq r+3$ by the following procedure.

First of all we define three rules for obtaining the vector

$$(Q_{j+1}^{j+1}, Q_{j+2}^{j+1}, Q_{j+3}^{j+1}, Q_{j+4}^{j+1})$$

from the vector

$$(Q_j^j, Q_{j+1}^j, Q_{j+2}^j, Q_{j+3}^j) :$$

rule $\mathfrak{R}_1$, rule $\mathfrak{R}_2$ and rule $\mathfrak{R}_3$. These rules correspond to different inequalities in Proposition 1.

Rule $\mathfrak{R}_1$:

$$\begin{cases} Q_{j+1}^{j+1} = Q_{j+1}^j, \\ Q_{j+2}^{j+1} = \max\{lQ_{j+1}^j, Q_{j+2}^j\}, \\ Q_{j+3}^{j+1} = \max\{lQ_{j+1}^j, Q_{j+3}^j\}, \\ Q_{j+4}^{j+1} = \max\{lQ_{j+1}^j, Q_{j+3}^j, Q_j^j + Q_{j+1}^j\}. \end{cases}$$

Rule $\mathfrak{R}_2$:

$$\begin{cases} Q_{j+1}^{j+1} = Q_{j+1}^j, \\ Q_{j+2}^{j+1} = Q_{j+2}^j, \\ Q_{j+3}^{j+1} = \max\{(2-l)Q_{j+1}^j + Q_j^j, Q_{j+3}^j\}, \\ Q_{j+4}^{j+1} = \max\{(2-l)Q_{j+1}^j + Q_j^j, Q_{j+3}^j, Q_j^j + Q_{j+1}^j\}. \end{cases}$$

Rule $\mathfrak{R}_3$:

$$\begin{cases} Q_{j+1}^{j+1} = Q_{j+1}^j, \\ Q_{j+2}^{j+1} = Q_{j+2}^j, \\ Q_{j+3}^{j+1} = Q_{j+3}^j, \\ Q_{j+4}^{j+1} = \max\{Q_{j+3}^j, Q_{j+2}^j + Q_{j+3}^j\}. \end{cases}$$

For a sequence $I = (i_0, \dots, i_{r-1})$ we take a sequence of rules $(\mathfrak{R}_{i_0}, \dots, \mathfrak{R}_{i_{r-1}})$ and construct a sequence $\{Q_j(I, m)\}, j = 0, \dots, r+3$ in the following way.

For $j = 0$ put

$$Q_0(m) = Q_0^0(m) = 1, \quad Q_t^0(m) = \theta_m^t, \quad t = 1, 2, 3.$$

For $j \geq 0$ given

$$(Q_j^j(m), Q_{j+1}^j(m), Q_{j+2}^j(m), Q_{j+3}^j(m))$$

we construct

$$(Q_{j+1}^{j+1}(m), Q_{j+2}^{j+1}(m), Q_{j+3}^{j+1}(m), Q_{j+4}^{j+1}(m))$$

by the rule $\mathfrak{R}_{i_j}$ with $l = l_m$.

Now we define $Q_j(I, m) = Q_j^j(m)$ for $j \leq r$ and $Q_{r+t, I}(m) = Q_{r+t}^r(m)$, $t = 1, 2, 3$.

The following proposition presents a result of computer calculation.

Proposition 2. *Let $m = 0, \dots, 6$. For any considered sequence of rules I and defined sequence $\{Q_k(I, m)\}$ one has*

$$(Q_{r+j}(I, m))^{\frac{1}{r+j}} \geq \theta_{m+1}, \quad j = 0, 1, 2, 3.$$

Remind that the increasing sequence of remainders of best approximations $\{q_k\}$ satisfies (2) and Proposition 1. When $l \in (0, 2)$ all coefficients in inequalities (6), (7), (8) are positive.

So we immediately deduce from Proposition 1 and Proposition 2 the following statement:

Proposition 3. *Suppose that*

$$q_{i+j} \geq \lambda \theta_m^j, \quad j = 0, 1, 2, 3, \quad \lambda > 0.$$

Then

$$q_{r+j} \geq \lambda \theta_{m+1}^{r+j}, \quad j = 0, 1, 2, 3.$$

From (2) it follows that for some positive λ one has $q_i \geq \lambda \theta_0^i$. By Proposition 3 we see that $q_{j+4r_0+2r_1} \geq \lambda \theta_7^{j+4r_0+2r_1}$ for any j . Theorem 3 is proved.

References

- [1] J.C.Lagarias, Best simultaneous diophantine approximation I. Growth rates of best approximation denominators. Trans. Am. Math. Soc., 272:545-554, 1980.
- [2] A.J.Brentjes, Multidimensional continued fraction algorithms, volume 145 of Mathematical Center Tracts. Mathematisch Centrum Amsterdam, 1982
- [3] M.V.Romanov, Simultaneous two-dimensional best Diophantine approximations in the Euclidean norm. Moscow Univ. Math. Bull. 61 (2006), no. 2, 34–37.

Author's address:

Dept. of Number Theory

Fac. Mathematics and Mechanics

Moscow State University

119992 Moscow

Russia

e-mail: zzremi@gmail.com